Allergy Action Plan



idents Name: D.O.B:		11 11 11 11 11 11 11 11 11 11 11 11 11
ALLERGY TO:		
Asthmatic		
SYMPTOMS:	Give checked medication as determined by Physician:	
If an exposure to the allergen has occurred, but there are NO symptoms:	☐ Antihistamine	
Mouth: Itching, tingling, or swelling of lips, tongue, mouth	□Antihistamine	□Epinephrine
Skin: Hives, itchy rash, swelling of the face or extremities	□Antihistamine	DEpinephrine
♦Gut: Nausea, abdominal cramps, vomiting, diarrhea	□Antihistamine	☐ Epinephrine
♦Throat: Tightening of throat, hoarseness, hacking cough	□Antihistamine	□Epinephrine
♦Lung: Shortness of breath, repetitive coughing, wheezing	□Antihistamine	☐ Epinephrine
♦Heart: Weak or thready pulse, low blood pressure, fainting, pale, blueness	□Antihistamine	☐ Epinephrine
Other symptoms:	□Antihistamine	□Epinephrine
If reaction is progressing, several of the above areas affected:	□Antihistamine	☐ Epinephrine
DOSAGE: Epinephrine: inject intramuscularly (see reverse side for instructions) □EpiPen® □EpiPen® Jr. □Twinject® 0.3 mg □Twinject® 0.15 mg		
Antihistamine: Medication/Dose/Route		
Other:		
Medication/Dose/Route IMPORTANT: Asthma inhalers and/or antihistamines cannot be depended anaphylaxis.	ed on to replace e	pinephrine in
EMERGENCY CALLS		
 Call 911. State that an allergic reaction has been treated, and additional epine Doctor:Phone Number: 		
	one Number:one Number:	
4. Other Emergency Contacts:		
* '		
Name: Phone Number:		
EVEN IF PARENT/GUARDIAN CANNOT BE REACHED, DO NOT HESITATE TO MEDICA MEDICAL FACILITY!		
Parent/Guardians Signature:	Date:	
Physicians Signature:		
Required	Date.	